

HADAMARD OPERATORS ON $\mathcal{D}'(\mathbb{R}^d)$

DIETMAR VOGT

BERGISCHE UNIVERSITÄT WUPPERTAL

FB MATH.-NAT., GAUSS-STR. 20,

D-42119 WUPPERTAL, GERMANY

E-MAIL: DVOGT@MATH.UNI-WUPPERTAL.DE

Dedicated to the memory of Paweł Domański

ABSTRACT. We study continuous linear operators on $\mathcal{D}'(\mathbb{R}^d)$ which admit all monomials as eigenvectors, that is, operators of Hadamard type. Such operators on $C^\infty(\mathbb{R}^d)$ and on the space $\mathcal{A}(\mathbb{R}^d)$ of real analytic functions on \mathbb{R}^d have been investigated by Domański, Langenbruch and the author. The situation in the present case, however, is quite different and also the characterization. An operator L on $\mathcal{D}'(\mathbb{R}^d)$ is of Hadamard type if there is a distribution T , the support of which has positive distance to all coordinate hyperplanes and which has a certain behaviour at infinity, such that $L(S) = S \star T$ for all $S \in \mathcal{D}'(\mathbb{R}^d)$. Here $(S \star T)\varphi = S_y(T_x\varphi(xy))$ for all $\varphi \in \mathcal{D}(\mathbb{R}^d)$. To describe the behaviour at infinity we introduce a class $\mathcal{O}'_H(\mathbb{R}^d)$ of distributions defined by the same conditions like in the description of class $\mathcal{O}'_C(\mathbb{R}^d)$ of Laurent Schwartz, but derivatives replaced with Euler derivatives.

In the present note we study Hadamard operators on $\mathcal{D}'(\mathbb{R}^d)$, that is, continuous linear operators on $\mathcal{D}'(\mathbb{R}^d)$ which admit all monomials as eigenvectors and we give a complete characterization. Such operators on $C^\infty(\mathbb{R}^d)$ have been studied and characterized in [9, 10], on $\mathcal{A}(\mathbb{R})$ in [1, 2, 3] and on $\mathcal{A}(\mathbb{R}^d)$ in [4]. There you find also references to the long history of such problems. Since it can be shown that Hadamard operators commute with dilations our problem is, by duality, closely related to the study of continuous linear operators in $\mathcal{D}(\mathbb{R}^d)$ which commute with dilations. They have the form $\varphi \mapsto T_x\varphi(xy)$ where T is a distribution. The class $\mathcal{D}'_H(\mathbb{R}^d)$ of distributions T such that $T_x\varphi(xy) \in \mathcal{D}(\mathbb{R}^d)$ for every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ is studied. These are distributions with positive distance to the coordinate hyperplanes and certain behaviour at infinity, similar to the class \mathcal{O}'_C of L. Schwartz of rapidly decreasing distributions. We define a class $\mathcal{O}'_H(\mathbb{R}^d)$ of distributions by the same conditions like in the description of class \mathcal{O}'_C in [7], but derivatives replaced with Euler derivatives. We denote by $\mathcal{M}(\mathbb{R}^d)$ the class of Hadamard

2010 *Mathematics Subject Classification.* Primary 46F10; Secondary 47B38, 46F05.

Key words and phrases. Operators on distributions, operators on test functions, monomials as eigenvectors, spaces of distributions.

operators in $\mathcal{D}'(\mathbb{R}^d)$ and for $x \in \mathbb{R}_*^d = (\mathbb{R} \setminus \{0\})^d$ we set $\sigma(x) = \prod_j \frac{x_j}{|x_j|}$, that is, the signum of x which, of course, is constant on each ‘quadrant’. Then our main result is (see Corollary 1.6, Theorem 2.10 and Theorem 4.2):

Main Theorem *$L \in \mathcal{M}(\mathbb{R}^d)$ if and only if there is a distribution $T \in \mathcal{O}'_H(\mathbb{R}^d)$, the support of which has positive distance to all coordinate hyperplanes, such that $L(S) = S \star T$ for all $S \in \mathcal{D}'(\mathbb{R}^d)$. Here $(S \star T)\varphi = S_y(T_x\varphi(xy))$ for all $\varphi \in \mathcal{D}(\mathbb{R}^d)$. The eigenvalues are $m_\alpha = T_x\left(\frac{\sigma(x)}{x^{\alpha+1}}\right)$.*

Moreover, we show that every Hadamard operator on $\mathcal{D}'(\mathbb{R}^d)$ maps $C^\infty(\mathbb{R}^d)$ to $C^\infty(\mathbb{R}^d)$, that is, defines a Hadamard operator on $C^\infty(\mathbb{R}^d)$. On the other hand not every Hadamard operator on $C^\infty(\mathbb{R}^d)$ can be extended to an operator on $\mathcal{D}'(\mathbb{R}^d)$.

We follow the notation in [9, 10]. The class of Hadamard operators in $C^\infty(\mathbb{R}^d)$, that is, of continuous linear operators which admit all monomials as eigenvectors is denoted by $M(\mathbb{R}^d)$. The Hadamard operators are given by distributions $T \in \mathcal{E}'(\mathbb{R}^d)$ by means of the formula $(M_T\varphi)(y) = T_x\varphi(xy)$. The dual $\mathcal{E}'(\mathbb{R}^d)$ is an algebra with respect to \star -convolution given by the formula $(T \star S)\varphi = T_x S_y\varphi(xy)$ where $xy = (x_1y_1, \dots, x_dy_d)$. $T \in \mathcal{E}'(\mathbb{R}^d)$ defines a \star -convolution operator $N_T : S \mapsto S \star T$ and $N_T = M_T^*$, that is, the dual operator of M_T .

Differential operators of the form $P(\theta)$ where P is a polynomial and $\theta_j = x_j\partial_j$ or, equivalently, of the form $\sum_\alpha c_\alpha x^\alpha \partial^\alpha$ are called Euler operators and θ_j is called an Euler derivative. On C^∞ these are the Hadamard operators M_T with $\text{supp } T = \{\mathbf{1}\}$ where $\mathbf{1} = (1, \dots, 1)$.

We use standard notation of Functional Analysis, in particular, of distribution theory. For unexplained notation we refer to [5], [6], [7], [8].

The author thanks Paweł Domański, who passed away too early, for useful discussions on the topic of this paper.

1. BASIC PROPERTIES

Definition 1. A map $L \in L(\mathcal{D}'(\mathbb{R}^d))$ is called a Hadamard operator if it admits all monomials as eigenvectors. The set of Hadamard operators we denote by $\mathcal{M}(\mathbb{R}^d)$.

Since the condition means that $L(x^\alpha) \in \text{span}\{x^\alpha\}$ for all $\alpha \in \mathbb{N}_0^d$ the set $\mathcal{M}(\mathbb{R}^d)$ is a closed subalgebra in $L_\sigma(\mathcal{D}'(\mathbb{R}^d))$ and therefore also in $L_b(\mathcal{D}'(\mathbb{R}^d))$. Here σ denotes the topology of pointwise convergence and b the topology of uniform convergence on bounded sets.

We define m_α by $L(x^\alpha) = m_\alpha x^\alpha$. Since the polynomials are dense in $\mathcal{D}'(\mathbb{R}^d)$ the operator $L \in \mathcal{M}(\mathbb{R}^d)$ is uniquely determined by the family m_α ,

$\alpha \in \mathbb{N}_0^d$, of eigenvalues. Clearly the set $\Lambda(\mathbb{R}^d)$ of eigenvalue families is an algebra and $L \rightarrow m_\alpha$ is an algebra isomorphism.

To study and characterize the Hadamard operators in $\mathcal{D}'(\mathbb{R}^d)$ we first need some preparations. We set $\mathbb{R}_* := \mathbb{R} \setminus \{0\}$. For $a \in \mathbb{R}_*^d$ we define the dilation operator $D_a \in L(\mathcal{D}'(\mathbb{R}^d))$ by

$$(D_a T)\varphi := T_x \left(\frac{\sigma(a)}{a_1 \dots a_d} \varphi\left(\frac{x}{a}\right) \right)$$

for $T \in \mathcal{D}'(\mathbb{R}^d)$ and $\varphi \in \mathcal{D}(\mathbb{R}^d)$. By direct verification we see that $D_a \xi^\alpha = a^\alpha \xi^\alpha$.

Lemma 1.1. *For $L \in \mathcal{M}(\mathbb{R}^d)$ and $a \in \mathbb{R}_*^d$ we have $L \circ D_a = D_a \circ L$.*

Proof. For any α we have $(L \circ D_a)\xi^\alpha = a^\alpha m_\alpha \xi^\alpha = (D_a \circ L)\xi^\alpha$. So the claim is shown for all polynomials and these are dense in $\mathcal{D}'(\mathbb{R}^d)$. \square

Notation. If $L \circ D_a = D_a \circ L$ for all a we say: L commutes with dilations.

By definition of D_a we obtain for the dual map $D_a^* \in L(\mathcal{D}(\mathbb{R}^d))$ of D_a :

Lemma 1.2. *For $a \in \mathbb{R}_*^d$ and $\varphi \in \mathcal{D}(\mathbb{R}^d)$ we have*

$$(D_a^* \varphi)(\xi) = \frac{\sigma(a)}{a_1 \dots a_d} \varphi\left(\frac{\xi}{a}\right).$$

If L commutes with dilations then $D_a^* \circ L^* = L^* \circ D_a^*$ for all $a \in \mathbb{R}_*^d$. For $\varphi \in \mathcal{D}(\mathbb{R}^d)$ we set $\psi = L^* \varphi$ and obtain

$$\frac{\sigma(a)}{a_1 \dots a_d} \psi\left(\frac{x}{a}\right) = L_\xi^* \left(\frac{\sigma(a)}{a_1 \dots a_d} \varphi\left(\frac{\xi}{a}\right) \right) [x].$$

For $\eta \in \mathbb{R}_*^d$ we set $a = 1/\eta$ and obtain $\psi(\eta x) = L_\xi^*(\varphi(\eta \xi))[x]$.

We have shown:

Lemma 1.3. *If $M = L^* \in L(\mathcal{D}(\mathbb{R}^d))$ and L commutes with dilations, then*

$$M_\xi(\varphi(\eta \xi))[x] = (M\varphi)(\eta x)$$

for all $\varphi \in \mathcal{D}(\mathbb{R}^d)$ and $\eta \in \mathbb{R}_*^d$.

For $\varphi \in \mathcal{D}(\mathbb{R}^d)$ we define now

$$T\varphi = (M\varphi)(\mathbf{1}) = (L\delta_1)(\varphi).$$

Then $T \in \mathcal{D}'(\mathbb{R}^d)$ and for all $\eta \in \mathbb{R}_*^d$ we have

$$(1) \quad (M\varphi)(\eta) = T_\xi \varphi(\eta \xi).$$

The problem with the right hand side of equation (1) is that for $\eta_j = 0$, in general, T cannot be applied to the non compact support function $\xi \rightarrow \varphi(\eta \xi)$. For T as above, however, the function $\eta \rightarrow T_\xi \varphi(\eta \xi)$, $\xi \in \mathbb{R}_*^d$, is the restriction of a function in $\mathcal{D}(\mathbb{R}^d)$.

Definition 2. By $\mathcal{D}'_H(\mathbb{R}^d)$ we denote the set of distributions $T \in \mathcal{D}'(\mathbb{R}^d)$ such that for every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the function $y \rightarrow T_\xi \varphi(\xi y)$, $y \in \mathbb{R}_*^d$, is the restriction of a function in $\mathcal{D}(\mathbb{R}^d)$.

This means that T must have the following properties:

- (*) For every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ there is $r > 0$ such that $T_\xi \varphi(\xi y) = 0$ for $y \in \mathbb{R}_*^d$ with $|y|_\infty > r$,
- (**) For every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the map $y \mapsto T_\xi \varphi(\xi y)$, $y \in \mathbb{R}_*^d$, extends to a function in $C^\infty(\mathbb{R}^d)$.

For $T \in \mathcal{D}'_H(\mathbb{R}^d)$ we denote by M_T the map which assigns to $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the continuous extension of $y \mapsto T_\xi \varphi(\xi y)$. From the closed graph theorem it follows easily:

Lemma 1.4. $M_T \in L(\mathcal{D}(\mathbb{R}^d))$ for every $T \in \mathcal{D}'_H(\mathbb{R}^d)$.

We obtain the following representation theorem:

Theorem 1.5. $L \in L(\mathcal{D}'(\mathbb{R}^d))$ commutes with dilations if and only if there is $T \in \mathcal{D}'_H(\mathbb{R}^d)$ such that $L(S) = S \star T$ for all $S \in \mathcal{D}'(\mathbb{R}^d)$, here $(S \star T)\varphi = S_y(T_x \varphi(xy))$ for $\varphi \in \mathcal{D}(\mathbb{R}^d)$. In this case $T = L(\delta_1)$.

Proof. The assertion can be written as $L = M_T^*$. If L commutes with translations then, by the above, $L^* = M_T$ for $T = L(\delta_1)$ and $T \in \mathcal{D}'_H(\mathbb{R}^d)$ by formula (1). This formula also implies the result. \square

Notation. For $T \in \mathcal{D}'_H(\mathbb{R}^d)$ we set $L_T(S) = S \star T$ for all $S \in \mathcal{D}'(\mathbb{R}^d)$.

Corollary 1.6. If $L \in \mathcal{M}(\mathbb{R}^d)$ then there is $T \in \mathcal{D}'_H(\mathbb{R}^d)$ such that $L = L_T$.

2. PROPERTIES OF $\mathcal{D}'_H(\mathbb{R}^d)$

First we will exploit the fact that $M_T \in L(\mathcal{D}(\mathbb{R}^d))$. For $\varepsilon > 0$ we set

$$W_\varepsilon = \{x \in \mathbb{R}^d : \min_j |x_j| \geq \varepsilon\}$$

and we will use the following notations:

For $r = (r_1, \dots, r_d)$, where all $r_j \geq 0$, we set $B_r = \{x \in \mathbb{R}^d : |x_j| \leq r_j \text{ for all } j\}$ and for $s \geq 0$ we set $B_s := B_{s\mathbf{1}} = \{x \in \mathbb{R}^d : |x|_\infty \leq s\}$.

For $r = (r_1, \dots, r_d) \in \mathbb{R}^d$ and $s \in \mathbb{R}$ we set $r+s = r+s\mathbf{1} = (r_1+s, \dots, r_d+s)$.

Lemma 2.1. 1. If $T \in \mathcal{D}'_H(\mathbb{R}^d)$ then there is $\varepsilon > 0$ such that $\text{supp } T \subset W_\varepsilon$.
 2. If $T \in \mathcal{D}'(\mathbb{R}^d)$ and there is $\varepsilon > 0$ such that $\text{supp } T \subset W_\varepsilon$ then T satisfies (*).

Proof. 1. If $T \in \mathcal{D}'_H(\mathbb{R}^d)$ there is $r > 0$ such that $\text{supp } M_T \varphi \subset B_r$ for any $\varphi \in \mathcal{D}(B_1)$ and this implies, that $T_\xi \varphi(\eta \xi) = 0$ for any $\varphi \in \mathcal{D}(B_1)$, $\eta \in \mathbb{R}_*^d$ and $|\eta|_\infty > r$.

We set $\varepsilon = 1/r$ and assume that $\text{supp } \varphi \cap W_\varepsilon = \emptyset$. That is,

$$\text{supp } \varphi \subset \bigcup_j \{x \in \mathbb{R}^d : |x_j| < \varepsilon\}.$$

Then we can write $\varphi = \sum_j \varphi_j$ with $\varphi_j \in \mathcal{D}(\{x \in \mathbb{R}^d : |x_j| < \varepsilon\})$.

We fix j and choose $\eta \in \mathbb{R}_*^d$ such that $\sup\{|x_\nu \eta_\nu| : x \in \text{supp } \varphi_j\} = 1$ for all ν . We set $\psi(\xi) = \varphi_j(\xi/\eta)$. Then $\text{supp } \psi \subset B_1$ and $|\eta|_\infty > r$ and therefore we have $T\varphi_j = T_\xi \psi(\eta \xi) = 0$.

Since this holds for every j , the proof of 1. is complete, the proof of 2. is obvious. \square

A special case is that of distributions with compact support.

Corollary 2.2. $\mathcal{E}'(\mathbb{R}^d) \cap \mathcal{D}'_H(\mathbb{R}^d) = \mathcal{E}'(\mathbb{R}_*^d)$.

Proof. This follows from Lemma 2.1 and the fact that (**) is fulfilled for distributions with compact support. \square

This will be used in Section 4 to handle the case of T with compact support.

After having settled property (*) of $T \in \mathcal{D}'_H(\mathbb{R}^d)$ we turn to property (**). It is quite restrictive.

Lemma 2.3. $\mathcal{D}'_H(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$.

Proof. Let $T \in \mathcal{D}'_H(\mathbb{R}^d)$, we may assume that $\text{supp } T \subset W_2$. For $k \in \mathbb{N}_0^d$ we set $|\varphi|_k = \|\varphi^{(k)}\|_{L_1}$ and remark that for every $\mathcal{D}(B_r)$ these norms are a fundamental system of seminorms. On $\mathcal{D}(B_1)$ the family of distributions $T_{(y)}\varphi := T_x \varphi(xy)$, $y \in \mathbb{R}_*^d \cap B_1$, is weakly bounded hence equicontinuous. This means that there is $k \in \mathbb{N}_0^d$ and $C > 0$ such that

$$|T_{(y)}\varphi| \leq C\|\varphi\|_k, \quad \varphi \in \mathcal{D}(B_1), |y|_\infty \leq 1.$$

For r with $r_j \geq 1$ for all j and $\varphi \in \mathcal{D}(B_r)$ we set $\psi(x) = \varphi(rx)$. Then $\psi \in \mathcal{D}(B_1)$ and therefore

$$|T\varphi| = |T_{(1/r)}\psi| \leq C\|\psi\|_k = Cr^k\|\varphi\|_k.$$

For every r there is $t_r \in L_\infty(B_r)$, $\|t_r\|_\infty \leq Cr^k$ such that $T\varphi = \int t_r(x) \varphi^{(k)}(x) dx$ for all $\varphi \in \mathcal{D}(B_r)$.

We restrict us now to $r \in \mathbb{N}^d$ and set $U_r = [r_1, r_1 + 1[\times \cdots \times [r_d, r_d + 1[$. Then $\bigcup_{r \in \mathbb{N}^d} U_r = W_1$. We choose $\chi \in \mathcal{D}(B_{1/2})$ with $\int \chi = 1$ and set $\chi_r(x) =$

$\int_{U_r} \chi(x - \xi) d\xi$. We obtain:

$$\begin{aligned}
T\varphi &= \sum_r T(\chi_r \varphi) = \sum_r \int t_{r+2}(x) (\chi_r \varphi)^{(k)}(x) dx \\
&= \sum_r \int t_{r+2}(x) \left(\sum_\nu \binom{k}{\nu} \chi_r^{(k-\nu)}(x) \varphi^{(\nu)}(x) \right) dx \\
&= \sum_\nu \binom{k}{\nu} \int \varphi^{(\nu)}(x) \left(\sum_r t_{r+2}(x) \chi_r^{(k-\nu)}(x) \right) dx \\
&= \sum_\nu \binom{k}{\nu} \int \varphi^{(\nu)}(x) \tau_\nu(x) dx.
\end{aligned}$$

We have to estimate the functions τ_ν . We set $\gamma(x) = \{r : x \in \prod_j [r_j - \frac{1}{2}, r_j + \frac{3}{2}]\}$

$$|\tau_\nu(x)| \leq \sum_{r \in \gamma(x)} |t_{r+2}(x)| |\chi_r^{(k-\nu)}(x)|.$$

For all 2^d elements $r \in \gamma(x)$ we have $|t_{r+2}(x)| \leq C(r+2)^k$ and $r \leq |x| + \frac{1}{2}$ (here $|x| = (|x_1|, \dots, |x_d|)$) and therefore $|t_{r+2}(x)| \leq C(|x| + 3)^k$. With a new constant C_ν we have

$$|\tau_\nu(x)| \leq C_\nu |x|^k.$$

This shows the result. \square

The necessary conditions we have found are far from being sufficient as the following example shows

Example 2.4. Let $d = 1$, we set $T\varphi = \int_1^\infty \varphi(x) dx$, then $T_x \varphi(xy) = \frac{1}{y} \int_y^\infty \varphi(x) dx$ for all $y > 0$ which, in general, is unbounded near 0. The distribution is $T \in \mathcal{S}'(\mathbb{R})$ and has support in W_1 .

The following definition is in analogy to a characterization of the space \mathcal{O}'_C of rapidly decreasing distributions in L. Schwartz [7], §5, p. 100.

Definition 3. $T \in \mathcal{O}'_H(\mathbb{R}^d)$ if for any k there are finitely many functions t_β such that $(1 + |x|^2)^{k/2} t_\beta \in L_\infty(\mathbb{R}^d)$ and such that $T = \sum_\beta \theta^\beta t_\beta$.

Proposition 2.5. If $T \in \mathcal{O}'_H(\mathbb{R}^d)$ then for every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the function $y \mapsto T_x(\varphi(xy))$, $y \in \mathbb{R}_*^d$, extends to a function in $C^\infty(\mathbb{R}^d)$.

Proof. We have to show that for every α the function $y \mapsto \partial_y^\alpha T_x(\varphi(xy))$, $y \in \mathbb{R}_*^d$, extends to a continuous function on \mathbb{R}^d .

By definition of $\mathcal{O}'_H(\mathbb{R}^d)$ we have to show our proposition only for $T = \theta^\beta t$ where $(1 + |x|^2)^{k/2} t \in L_\infty(\mathbb{R}^d)$ and k is suitably chosen. Since the formal

adjoint $(\theta^*)^\beta$ of the Euler-operator θ^β is again an Euler-operator we have

$$\int (\theta^\beta t)(x) \varphi(x) dx = \int t(x) ((\theta^*)^\beta \varphi)(x) dx = \sum_{\nu} c_{\nu} \int t(x) x^{\nu} \varphi^{(\nu)}(x) dx$$

where the sum is finite with suitable c_{ν} . Therefore it is enough to study the function

$$(2) \quad F(y) := \int t(x) (xy)^{\nu} \varphi^{(\nu)}(xy) dx, \quad y \in \mathbb{R}_*^d.$$

We have to show that all limits $\lim_{y \rightarrow y_0} F^{(\alpha)}(y)$, $y_0 \in \mathbb{R}^d$, $\alpha \in \mathbb{N}_0^d$ exist.

$$(3) \quad F^{(\alpha)}(y) = \int t(x) x^{\alpha} (x^{\nu} \varphi^{(\nu)}(x))^{(\alpha)} [xy] dx \\ = \sum_{\alpha - \nu \leq \gamma \leq \alpha} c_{\gamma} \int t(x) x^{\alpha} (xy)^{\nu - \alpha + \gamma} \varphi^{(\nu + \gamma)}(xy) dx.$$

If k was chosen so large that $t(x) x^{\alpha} \in L_1(\mathbb{R}^d)$ then the limits exist as requested. \square

Therefore we have shown:

Theorem 2.6. *If $T \in \mathcal{O}'_H(\mathbb{R}^d)$ and $\text{supp } T \subset W_{\varepsilon}$ for some $\varepsilon > 0$, then $T \in \mathcal{D}'_H(\mathbb{R}^d)$.*

To show the inverse we give a description of $\mathcal{D}'_H(\mathbb{R}^d)$ in terms of the regularizations of a distribution.

Theorem 2.7. *T satisfies $(**)$ if and only if $T * \chi$ satisfies $(**)$ for all $\chi \in L_1(B_1)$.*

*In this case the set of maps $\{\varphi \mapsto (T * \chi)_x \varphi(x \cdot) : \chi \in L_1(B_1), \|\chi\|_{L_1} \leq 1\}$ is equicontinuous in $L(\mathcal{D}(\mathbb{R}^d), C^\infty(\mathbb{R}^d))$.*

Proof. We assume that T satisfies $(**)$ and want to show that the same holds for $T * \chi$. We need some preparation. We set:

$$\psi(\xi, y_2) = \int \chi(x_2) \varphi(\xi + x_2 y_2) dx_2$$

and remark that for $\varphi \in \mathcal{D}(B_r)$ and $\chi \in L_1(B_1)$ we have $\psi(\cdot, y_2) \in \mathcal{D}(B_{r+|y_2|})$. This implies that the map which assigns to every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the function $y_2 \mapsto \psi(\cdot, y_2)$ is a continuous linear map $\Phi : \mathcal{D}(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}^d, \mathcal{D}(\mathbb{R}^d))$.

We set

$$F(y_1, y_2) = T_{x_1} \psi(x_1 y_1, y_2) \in C^\infty(\mathbb{R}_*^d \times \mathbb{R}^d).$$

By assumption it extends to a function \hat{F} on $\mathbb{R}^d \times \mathbb{R}^d$ such that $\hat{F}(\cdot, y_2)$ is in $C^\infty(\mathbb{R}^d)$ for every $y_2 \in \mathbb{R}^d$.

The map which assigns to every $g \in \mathcal{D}(\mathbb{R}^d)$ the continuous extension of $T_\xi g(\xi y_1)$ defines a continuous linear map $\Psi : \mathcal{D}(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}^d)$. We obtain

$$\hat{F}(y_1, y_2) = \Psi\{\Phi(\varphi)[y_2]\}[y_1] \in C^\infty(\mathbb{R}^d, C^\infty(\mathbb{R}^d)) = C^\infty(\mathbb{R}^d \times \mathbb{R}^d).$$

Therefore $\hat{F}(y, y) \in C^\infty(\mathbb{R}^d)$ and it is the extension of

$$F(y, y) = T_{x_1} \int \chi(x_2) \varphi((x_1 + x_2)y) dx_2 = (T * \chi)_x \varphi(xy).$$

This proves one direction of the Theorem, it remains to show the other implication. We will use the idea of proof in [7, §7, Théorème XX].

First we show the additional assertion of the theorem. We assume that $T * \chi$ satisfies $(**)$ for all $\chi \in L_1(B_1)$. We consider the map $L_1(B_1) \rightarrow L(\mathcal{D}(\mathbb{R}^d), C^\infty(\mathbb{R}^d))$ defined by $\chi \mapsto [\varphi \mapsto (T * \chi)_x \varphi(x \cdot)]$ (cf. Lemma 1.4). If $\|\chi\| \rightarrow 0$ and $[\varphi \mapsto (T * \chi)_x \varphi(x \cdot)] \rightarrow A$ in $L(\mathcal{D}(\mathbb{R}^d), C^\infty(\mathbb{R}^d))$ then for fixed $y \in \mathbb{R}_*^d$ and all φ we have $(T * \chi)_x \varphi(xy) \rightarrow 0$ and therefore $(A\chi)\varphi = 0$ on \mathbb{R}_*^d , hence on \mathbb{R}^d . So the map $L_1(B_1) \rightarrow L(\mathcal{D}(\mathbb{R}^d), C^\infty(\mathbb{R}^d))$ has closed graph and, by de Wilde's Theorem, is continuous. This shows the assertion.

We fix $\chi \in \mathcal{D}(\mathbb{R}^d)$ with $\int |\chi| = 1$. Then for every $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the function $(T * \chi)_x \varphi(xy)$, $y \in \mathbb{R}_*^d$, extends to a function in $C^\infty(\mathbb{R}^d)$ and the set $\{F_\chi : \varphi \mapsto (T * \chi)_x \varphi(xy) : \chi \in \mathcal{D}(\mathbb{R}^d), \chi \geq 0, \int \chi = 1\}$ is equicontinuous. Therefore it is relatively compact in $L(\mathcal{D}(B_R), C^\infty(\mathbb{R}^d))$. We fix χ and set $\chi_\varepsilon(x) = \varepsilon^{-d} \chi(x/\varepsilon)$ for $\varepsilon > 0$. Then there is a sequence $\varepsilon_n \downarrow 0$ such that $F_{\chi_{\varepsilon_n}}$ converges to some $F \in L(\mathcal{D}(B_R), C^\infty(\mathbb{R}^d))$. Since

$$F_{\chi_{\varepsilon_n}} = T_\xi \left(\int \chi_{\varepsilon_n}(x) \varphi((x + \xi)y) dx \right) \rightarrow T_\xi \varphi(\xi y)$$

for every $y \in \mathbb{R}_*^d$ we see, that $T_\xi \varphi(\xi y)$ extends to a function in $C^\infty(\mathbb{R}^d)$. \square

Now we can show the inverse of Theorem 2.6.

Theorem 2.8. *If $T \in \mathcal{D}'_H(\mathbb{R}^d)$ then for every $\beta > 0$ there is a function t_β such that $x^\beta t_\beta$ is bounded and an Euler operator $P(\theta)$ such that $T = P(\theta)t_\beta$. In particular $T \in \mathcal{O}'_H(\mathbb{R}^d)$.*

Proof. Let $T \in \mathcal{D}'_H(\mathbb{R}^d)$ with $\text{supp } T \subset W_{2\varepsilon_0}$. Let $\chi \in \mathcal{D}(\mathbb{R}^d)$ with $\chi \geq 0$, $\int \chi = 1$ and $\text{supp } \chi \subset B_{\varepsilon_0}$. Then for every $\varphi \in \mathcal{D}(B_1)$ the function $(T * \chi)_x \varphi(xy)$, $y \in \mathbb{R}_*^d$, extends to a function in $\mathcal{D}(\mathbb{R}^d)$. We set $\tau = T * \chi$.

We set for $y \in \mathbb{R}_*^d$ and $\varphi \in \mathcal{D}(\mathbb{R}^d)$:

$$F(y) = \int \tau(x) \varphi(xy) dx.$$

Then $F \in C^\infty(\mathbb{R}_*^d)$, $F(y) = 0$ for $|y|_\infty > 1/\varepsilon_0$. For $\beta \in \mathbb{N}_0$, we have

$$F^{(\beta)}(y) = \int \tau(x) x^\beta \varphi^{(\beta)}(xy) dx.$$

By Theorem 2.7 on $\mathcal{D}(B_1)$ the set of distributions $\varphi \rightarrow F^{(\beta)}(y)$, $y \in \mathbb{R}_*^d$, χ as above, is equicontinuous. Hence there is p such that all these distributions extend to $\mathcal{D}^{|\beta|+p}(B_1)$ and the set of these distributions is bounded in $\mathcal{D}^{|\beta|+p}(B_1)'$.

For $\alpha \in \mathbb{N}_0$ we choose $\varphi_\alpha \in \mathcal{D}^\alpha[0, +1]$ such that $\varphi_\alpha \in C^\infty(\mathbb{R} \setminus \{0, 1\})$ and

$$\varphi_\alpha(x) = x^{\alpha+1} \frac{(x-1)^{\alpha+1}}{(2\alpha+2)!} \text{ for } 0 \leq x \leq 1.$$

We define for $\alpha \in \mathbb{N}_0^d$ such that $\mathcal{D}^{|\alpha|}(B_1) \subset \mathcal{D}^{|\beta|+p}(B_1)$

$$\varphi_\alpha(x) = \prod_{j=1}^d \varphi_{\alpha_j}(x_j)$$

and consider $F(y) = \int \tau(x) \varphi_\alpha(xy) dx$. Then we have for $y \in (0, +\infty)^d$, setting $\alpha + 1 = (\alpha_1 + 1, \dots, \alpha_d + 1)$, etc.

$$F^{(2\alpha+2)}(y) = \int_{[0, 1/y]^d} \tau(x) x^{2\alpha+2} dx$$

and therefore

$$F^{(2\alpha+3)}(y) = -\frac{1}{y^{2\alpha+4}} \tau\left(\frac{1}{y}\right).$$

With an analogous argument for the other ‘quadrants’ we get for general $y \in \mathbb{R}_*^d$

$$F^{(2\alpha+3)}(y) = -\frac{\sigma(y)}{y^{2\alpha+4}} \tau\left(\frac{1}{y}\right).$$

We set $G(x) = \sigma(x) F^{(\beta)}(\frac{1}{x})$. Then G is a bounded function on \mathbb{R}_*^d with a bound independent of χ and $\sigma(x) F^{(\beta)}(x) = G(\frac{1}{x})$. We calculate the derivatives. With certain coefficients c_ν we have:

$$\sigma(x) F^{(\beta+q)}(x) = \sum_{\nu}' c_\nu \frac{1}{x^{\nu+q}} G^{(\nu)}\left(\frac{1}{x}\right)$$

where \sum_{ν}' runs over $1 \leq \nu_j \leq q$. We choose $q = 2\alpha + 3 - \beta$. Replacing x with $1/x$ and c_ν with $-c_\nu$ we obtain:

$$(4) \quad x^{\beta+1} \tau(x) = \sum_{\nu}' c_\nu x^\nu G^{(\nu)}(x).$$

We analyze now functions of type $x^{m+\nu} G^{(\nu)}(x)$ and claim

Lemma 2.9. *Any such function can be written as a linear combination of the functions $(x^{m+j} G(x))^{(j)}$, $0 \leq j_i \leq \nu_i$ for all i .*

Proof. For any $m \in \mathbb{Z}$ there are constants $c_{m,\nu,j}$ such that

$$(x^{m+\nu} G(x))^{(\nu)} = \sum_{j=0}^{\nu} c_{m,\nu,j} y^{m+\nu-j} G^{(\nu-j)}(x)$$

and therefore, since $c_{m,\nu,0} = 1$,

$$x^{m+\nu}G^{(\nu)}(x) = (x^{m+\nu}G(x))^{(\nu)} - \sum_j' c_{m,\nu,j} x^{m+\nu-j}G^{(\nu-j)}(x).$$

where \sum_j' runs over all j with $j \neq 0$ and $0 \leq j_i \leq \nu_i$ for all i . Induction over $|\nu|$ yields the result. \square

End of the proof of Theorem 2.8: From Lemma 2.9 and equation (4) we obtain with new constants c_j :

$$\tau(x) = \sum_{|j|_\infty \leq p+2} c_j (x^j (x^{-\beta-1}G(x)))^{(j)}.$$

We fix now $\chi \in \mathcal{D}(\mathbb{R}^d)$ with $\chi \geq 0$ and $\int \chi = 1$. For $\varepsilon > 0$ we set $\chi_\varepsilon = \varepsilon^{-d}\chi(x/\varepsilon)$, $\tau_\varepsilon = T * \chi_\varepsilon$ and G_ε the corresponding function. Then we have

$$(5) \quad (T * \chi_\varepsilon)\varphi = \int \tau_\varepsilon(x)\varphi(x)dx = \sum_{|j|_\infty \leq p+2} c_j \int x^{-\beta-1}G_\varepsilon(x)x^j\varphi^{(j)}(x)dx.$$

We have $\lim_{\varepsilon \rightarrow 0}(T * \chi_\varepsilon)\varphi = T\varphi$. On the other hand $\{G_\varepsilon : \varepsilon > 0\}$ is bounded in $L_\infty(\mathbb{R}^d) = L_1(\mathbb{R}^d)'$. Therefore there is $G \in L_\infty(\mathbb{R}^d)$ and a sequence G_{ε_n} which converges to G in the weak*-topology with respect to $L_1(\mathbb{R}^d)$.

From equation (5) then follows:

$$\begin{aligned} T\varphi &= \sum_{|j|_\infty \leq p+2} c_j \int x^{-\beta-1}G(x)x^j\varphi^{(j)}(x)dx = \int (x^{-\beta-1}G(x))(P(\theta)\varphi)(x)dx \\ &= \int (P(\theta)^*(x^{-\beta-1}G(x))\varphi(x)dx. \end{aligned}$$

Putting $t_\beta := x^{-\beta-1}G(x)$ we have shown: For every β there is a function t_β such that $x^\beta t_\beta \in L_\infty(\mathbb{R}^d)$ and an Euler operator $Q(\theta) := P(\theta)^*$ such that $T = Q(\theta)t_\beta$. This completes the proof. \square

From Theorems 2.6 and 2.8 we obtain one of the main results of this paper:

Theorem 2.10. $\mathcal{D}'_H(\mathbb{R}^d) = \{T \in \mathcal{O}'_H(\mathbb{R}^d) : \text{supp } T \subset W_\varepsilon \text{ for some } \varepsilon > 0\}.$

3. THE SPACE $\mathcal{O}'_H(\mathbb{R}^d)$

We recall the definition of the space $\mathcal{O}'_H(\mathbb{R}^d)$:

$T \in \mathcal{O}'_H(\mathbb{R}^d)$ if for any k there are finitely many functions t_β such that $(1 + |x|^2)^{k/2}t_\beta \in L_\infty(\mathbb{R}^d)$ and such that $T = \sum_\beta \theta^\beta t_\beta$.

The space $\mathcal{O}'_C(\mathbb{R}^d)$ of L. Schwartz may be defined by any of the following equivalent properties (see [7, §5, Théorème IX]). Let $T \in \mathcal{D}'(\mathbb{R}^d)$, then $T \in \mathcal{O}'_C(\mathbb{R}^d)$ if and only if 1. or 2.:

- (1) For any k there are finitely many functions t_β such that $(1+|x|^2)^{k/2}t_\beta \in L_\infty(\mathbb{R}^d)$ and such that $T = \sum_\beta \partial^\beta t_\beta$.
- (2) For any $\chi \in \mathcal{D}(\mathbb{R}^d)$, $T*\chi$ is a rapidly decreasing continuous function.

Without proof we admit.

Lemma 3.1. $\mathcal{E}'(\mathbb{R}^d) \subset \mathcal{O}'_C(\mathbb{R}^d) \cap \mathcal{O}'_H(\mathbb{R}^d)$.

And this implies:

Lemma 3.2. If $T|_{\mathbb{R}^d \setminus B_R} = S|_{\mathbb{R}^d \setminus B_R}$ for some $R > 0$ and $S \in \mathcal{O}'_H(\mathbb{R}^d)$ or $S \in \mathcal{O}'_C(\mathbb{R}^d)$ then $T \in \mathcal{O}'_H(\mathbb{R}^d)$ or $T \in \mathcal{O}'_C(\mathbb{R}^d)$, respectively.

Proof. $T = S + (T - S)$ and $T - S \in \mathcal{E}'(\mathbb{R}^d)$. □

Proposition 3.3. 1. $\mathcal{O}'_C(\mathbb{R}) \subset \mathcal{O}'_H(\mathbb{R})$.

2. If $\text{supp } T \subset W_\varepsilon$ for some $\varepsilon > 0$ and $T \in \mathcal{O}'_C(\mathbb{R}^d)$ then $T \in \mathcal{O}'_H(\mathbb{R}^d)$.

Proof. 2. It is enough to show the claim for $T = t_\beta^{(\beta)}, (1+|x|^2)^{k/2}t_\beta$ bounded. Set $\tau_\beta = \frac{1}{x^\beta}t_\beta$. Then τ_β is decreasing even faster and $(x^\beta \tau_\beta)^{(\beta)} = t_\beta^{(\beta)}$. Since $f \mapsto (x^\beta f)^{(\beta)}$ is an Euler operator the proof of 2. is complete.

1. Choose $\chi \in \mathcal{D}[-1, +1]$, $\chi \equiv 1$ in a neighborhood of 0. For $T \in \mathcal{O}'_C(\mathbb{R})$ set $S = (1 - \chi)T$. Then $\text{supp } S \subset W_\varepsilon$ for some $\varepsilon > 0$ and, due to Lemma 3.2, $S \in \mathcal{O}'_C(\mathbb{R})$. By part 2. of this Proposition we have $S \in \mathcal{O}'_H(\mathbb{R})$ and therefore, again by Lemma 3.2, $T \in \mathcal{O}'_H(\mathbb{R})$. □

The space $\mathcal{O}'_C(\mathbb{R})$ is a proper subspace of $\mathcal{O}'_H(\mathbb{R})$, as the following example shows.

Example 3.4. If $T = e^{-ix}$, that is, $T\varphi = \int e^{-ix}\varphi(x)dx$, then

1. $T \notin \mathcal{O}'_C(\mathbb{R})$, 2. $T \in \mathcal{O}'_H(\mathbb{R})$.

Proof. 1. Let $\chi \in \mathcal{D}(\mathbb{R})$, then $(T * \chi)(x) = \int e^{-i\xi}\chi(x - \xi)d\xi = \hat{\chi}(-1)e^{-ix}$. By the second definition of $\mathcal{O}'_C(\mathbb{R})$ (see above) $T \notin \mathcal{O}'_C(\mathbb{R})$.

2. To show that $T \in \mathcal{O}'_H(\mathbb{R})$ we choose $\chi \in \mathcal{D}[-1, +1]$, $\chi \equiv 1$ in a neighborhood of 0. For given k we set $t_k(x) = \frac{i^k}{x^k}(1 - \chi(x))e^{-ix}$. Then $(1+x^2)^{k/2}t_k(x)$ is bounded and $(x^k t_k(x))^{(k)} = (i^k(1 - \chi(x))e^{-ix})^{(k)} = e^{-ix} + g(x)$ where g has compact support. Hence $T = (x^k t_k(x))^{(k)} - g$ where g has compact support. This shows the result like above. □

By Proposition 2.5 we know now that for T as in Example 3.4 the function $T_x\varphi(xy)$, $y \in \mathbb{R}_*$, extends to a C^∞ -function on \mathbb{R} . For this example we can make it explicit, even for higher dimensions, setting $e^{-ix} = e^{-i(x_1 + \dots + x_d)}$.

For $\varphi \in \mathcal{D}(\mathbb{R}^d)$ and $y \in \mathbb{R}_*^d$ we set $F(y) = T_x \varphi(xy)$. We obtain for $y \in \mathbb{R}_*^d$

$$\begin{aligned} \partial^\alpha F(y) &= \int e^{-ix} x^\alpha \varphi^{(\alpha)}(xy) dx = \frac{\sigma(y)}{y^{\alpha+1}} \int e^{-i(x/y)} x^\alpha \varphi^{(\alpha)}(x) dx \\ &= \frac{\sigma(y)}{y^{\alpha+1}} \widehat{x^\alpha \varphi^{(\alpha)}}\left(\frac{1}{y}\right). \end{aligned}$$

Since $\widehat{x^\alpha \varphi^{(\alpha)}} \in \mathcal{S}(\mathbb{R}^d)$ we obtain $\lim_{y \rightarrow y_0} F^{(\alpha)}(y) = 0$ for every $y_0 \in \mathbb{R}^d \setminus \mathbb{R}_*^d$. That means, if we denote the extended function again by F , that $F^{(\alpha)}(y) = 0$ on all coordinate hyperplanes and for all α .

Returning to the one-dimensional case we present another example which we take from [7, §5, p. 100, (VII,5;1)].

Example 3.5. If $T = e^{i\pi x^2}$ then $T \in \mathcal{O}'_C(\mathbb{R})$ and therefore $T \in \mathcal{O}'_H(\mathbb{R})$. The function $e^{i\pi x^2}$ is bounded, but its derivatives are not.

4. EIGENVALUES

In this section we study Hadamard operators in terms of the representing distribution in $\mathcal{D}'_H(\mathbb{R}^d)$. A special case are the distributions in $\mathcal{D}'_H(\mathbb{R}^d)$ with compact support.

In consequence of Lemma 1.1, Theorem 1.5 and Corollary 2.2 we obtain:

Theorem 4.1. *For $T \in \mathcal{D}'(\mathbb{R}^d)$ the following are equivalent:*

1. $T \in \mathcal{E}'(\mathbb{R}^d)$ and the \star -homomorphism N_T can be extended to a map in $\mathcal{M}(\mathbb{R}^d)$.
2. $T \in \mathcal{D}'_H(\mathbb{R}^d)$ and $L_T \mathcal{E}'(\mathbb{R}^d) \subset \mathcal{E}'(\mathbb{R}^d)$.
3. $T \in \mathcal{E}'(\mathbb{R}_*^d)$.

In this case $m_\alpha = T_x \left(\frac{\sigma(x)}{x^{\alpha+1}} \right)$.

Proof. 1. \Rightarrow 2.: By assumption there is $L \in \mathcal{M}(\mathbb{R}^d)$ such that $L(S) = N_T(S) = S \star T$ for all $S \in \mathcal{E}'(\mathbb{R}^d)$. Then $T = N_T(\delta_1) = L(\delta_1) \in \mathcal{D}'_H(\mathbb{R}^d)$. By definition $N_T(S) = S \star T = L_T(S)$ for $S \in \mathcal{E}'(\mathbb{R}^d)$.

2. \Rightarrow 3.: By assumption $T = L_T(\delta_1) \in \mathcal{E}'(\mathbb{R}^d)$. Hence $T \in \mathcal{E}'(\mathbb{R}^d) \cap \mathcal{D}'_H(\mathbb{R}^d) = \mathcal{E}'(\mathbb{R}_*^d)$, by Corollary 2.2.

3. \Rightarrow 1. By Corollary 2.2 $T \in \mathcal{E}'(\mathbb{R}_*^d) \subset \mathcal{D}'_H(\mathbb{R}^d)$ and we have $L_T(S) = S \star T = N_T(S) \in \mathcal{E}'(\mathbb{R}^d)$ for $S \in \mathcal{E}'(\mathbb{R}^d)$.

Moreover

$$\begin{aligned} L_T(\xi^\alpha)[\varphi] &= \int \xi^\alpha (T_x \varphi(x\xi)) d\xi = T_x \left(\frac{1}{x^\alpha} \int (x\xi)^\alpha \varphi(x\xi) d\xi \right) \\ &= T_x \left(\frac{\sigma(x)}{x^{\alpha+1}} \int \eta^\alpha \varphi(\eta) d\eta \right) \\ &= \int (m_\alpha \eta^\alpha) \varphi(\eta) d\eta. \end{aligned}$$

This shows that $L_T(\xi^\alpha) = m_\alpha \xi^\alpha$ with $m_\alpha = T_x\left(\frac{\sigma(x)}{x^{\alpha+1}}\right)$. \square

Notice that the interchange of T and the integral is clear since $T \in \mathcal{E}'(\mathbb{R}^d)$. For T with non-compact support this is more complicated. We further study operators represented by distributions in $\mathcal{D}'_H(\mathbb{R}^d)$. By Theorems 1.5 and 2.10 we know that they define operators $L_T \in L(\mathcal{D}'(\mathbb{R}^d))$ commuting with dilations. We show that these are, in fact, Hadamard operators.

Theorem 4.2. *If $T \in \mathcal{D}'_H(\mathbb{R}^d)$, then L_T is a Hadamard operator with eigenvalues $m_\alpha = T_x\left(\frac{\sigma(x)}{x^{\alpha+1}}\right)$.*

Proof. It remains to show that L_T admits all monomials as eigenvectors. It is sufficient to assume that $T = (-1)^{|k|}(x^k \tau)^{(k)}$, where $\tau \in L_1(\mathbb{R}^d)$ and $k \in \mathbb{N}_0^d$. Then the function

$$f(x, \xi) = \xi^\alpha \tau(x) \xi^k \varphi^{(k)}(x\xi)$$

is in $L_1(\mathbb{R}^d \times \mathbb{R}^d)$. To see this let $\text{supp } \varphi \subset B_R$. Then $f(x, \xi) \neq 0$ only if $|x_j| \geq \varepsilon$ and $|x_j \xi_j| \leq R$ for all j , hence only for $|\xi_j| \leq R/\varepsilon$ for all j .

$$\begin{aligned} \int \xi^\alpha T_x \varphi(x\xi) d\xi &= \int \xi^\alpha \left(\int \tau(x) (x\xi)^k \varphi^{(k)}(x\xi) dx \right) d\xi \\ &= \int \tau(x) \frac{1}{x^\alpha} \left(\int (x\xi)^{\alpha+k} \varphi^{(k)}(x\xi) d\xi \right) dx \\ &= \int \tau(x) \frac{\sigma(x)}{x^{\alpha+1}} \left(\int \eta^{\alpha+k} \varphi^{(k)}(\eta) d\eta \right) dx \\ &= \int \tau(x) \frac{\sigma(x)}{x^{\alpha+1}} (-1)^{|k|} \frac{(\alpha+k)!}{\alpha!} \left(\int \eta^\alpha \varphi(\eta) d\eta \right) dx. \end{aligned}$$

Therefore

$$\int \xi^\alpha T_x \varphi(x\xi) d\xi = \int (m_\alpha \eta^\alpha) \varphi(\eta) d\eta$$

where

$$m_\alpha = (-1)^{|k|} \frac{(\alpha+k)!}{\alpha!} \int \tau(x) \frac{\sigma(x)}{x^{\alpha+1}} dx.$$

Taking into account that for x with $\min_j |x_j| > 0$

$$x^k \left(\frac{\sigma(x)}{x^{\alpha+1}} \right)^{(k)} = (-1)^{|k|} \frac{(\alpha+k)!}{\alpha!} \frac{\sigma(x)}{x^{\alpha+1}}$$

we finally obtain

$$m_\alpha = T\left(\frac{\sigma(x)}{x^{\alpha+1}}\right)$$

which proves the result. \square

REMARK: We needed in the proof only very weak assumptions on T . Under these $\varphi \mapsto T_x \varphi(xy)$ might not send $\mathcal{D}(\mathbb{R}^d)$ into $\mathcal{D}(\mathbb{R}^d)$ hence its adjoint sends something much smaller to $\mathcal{D}'(\mathbb{R}^d)$, but it sends ξ^α to $m_\alpha \xi^\alpha$ for all $\alpha \in \mathbb{N}_0^d$.

5. HADAMARD OPERATORS IN $\mathcal{D}'(\mathbb{R}^d)$ AND IN $C^\infty(\mathbb{R}^d)$

We study now the problem when a Hadamard operator M on $C^\infty(\mathbb{R}^d)$ extends to an operator on $\mathcal{D}'(\mathbb{R}^d)$ and, on the other side, when an operator $L \in \mathcal{M}(\mathbb{R}^d)$ leaves $C^\infty(\mathbb{R}^d)$ invariant, that is, $LC^\infty(\mathbb{R}^d) \subset C^\infty(\mathbb{R}^d)$.

We start with the latter question, which has a rather straightforward answer.

Theorem 5.1. *If $T \in \mathcal{D}'_H(\mathbb{R}^d)$ then $L_T \in \mathcal{M}(\mathbb{R}^d)$ and $L_T(C^\infty(\mathbb{R}^d)) \subset C^\infty(\mathbb{R}^d)$.*

Proof. We may assume that $T = P(\theta)t$ where $t \in L_1(\mathbb{R}^d)$ with $\text{supp } t \subset W_\varepsilon$ for some $\varepsilon > 0$ and $P(\theta)$ is an Euler operator. Let $P^*(\theta)$ denote the formal adjoint of $P(\theta)$ which is again an Euler operator. Then for $f \in C^\infty(\mathbb{R}^d)$, $\varphi \in \mathcal{D}(\mathbb{R}^d)$ the function

$$f(y)t(x)P^*(\theta)_x\varphi(xy), \quad x, y \in \mathbb{R}^d$$

is in $L_1(\mathbb{R}^d \times \mathbb{R}^d)$. We obtain, using that Euler operators commute with dilations:

$$\begin{aligned} (L_T f)\varphi &= \int f(y) \left(\int t(x) P^*(\theta)_x \varphi(xy) dx \right) dy \\ &= \int \int f(y) t(x) (P^*(\theta)\varphi)(xy) dx dy. \end{aligned}$$

We apply the substitution $xy = \eta$, $y = \xi\eta$, its Jacobian determinant is $\frac{(-1)^d}{\xi_1 \cdots \xi_d}$.

$$(L_T f)\varphi = \int \left(\int f(\xi\eta) t\left(\frac{1}{\xi}\right) \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d} d\xi \right) (P^*(\theta)\varphi)(\eta) d\eta.$$

We have shown that on $C^\infty(\mathbb{R}^d)$ we have $L_T = P(\theta) \circ M_{T^\#}$ where

$$T^\# = t\left(\frac{1}{\xi}\right) \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d}.$$

Notice that $t\left(\frac{1}{\xi}\right) \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d}$ is an $L_1(\mathbb{R}^d)$ -function with compact support, hence $T^\# \in \mathcal{E}'(\mathbb{R}^d)$. \square

REMARK: Like in Theorem 4.2 we needed much weaker assumptions than $T \in \mathcal{O}'_H(\mathbb{R}^d)$, cf. the remark at the end of Section 4. This corresponds to the fact that not all Hadamard operators in $C^\infty(\mathbb{R}^d)$ extend to operators in $\mathcal{M}(\mathbb{R}^d)$ as the following proposition shows:

Proposition 5.2. *If $\text{supp } T = \{0\}$ and $T \neq 0$ then M_T cannot be extended to a map in $\mathcal{M}(\mathbb{R}^d)$*

Proof. If $T = \sum_{\beta} c_{\beta} \delta^{(\beta)}$ then $M_T(f)[x] = \sum_{\beta} c_{\beta} (-1)^{|\beta|} f^{(\beta)}(0) x^{\beta}$, hence $R(M_T) \subset E = \text{span}\{x^{\beta} : \beta \in e\}$ where e is a finite set. Assume $L \in \mathcal{M}(\mathbb{R}^d)$ and $L|_{C^{\infty}(\mathbb{R}^d)} = M_T$. Since E is closed in $\mathcal{D}'(\mathbb{R}^d)$ and $C^{\infty}(\mathbb{R}^d)$ is dense in $\mathcal{D}'(\mathbb{R}^d)$ we have $R(L) \subset E$ and this implies that $L(S) = \sum_{\beta} S(\varphi_{\beta}) x^{\beta}$ for all $S \in \mathcal{D}'(\mathbb{R}^d)$ where $\varphi_{\beta} \in \mathcal{D}(\mathbb{R}^d)$ for all $\beta \in e$. This implies $c_{\beta} (-1)^{\beta} f^{(\beta)}(0) = \int f(\xi) \varphi_{\beta}(\xi) d\xi$ for all $\beta \in e$ and $f \in C^{\infty}(\mathbb{R}^d)$ which is possible only if $c_{\beta} = 0$ for all $\beta \in e$, that is, $T = 0$. \square

To study the problem which Hadamard operators on $C^{\infty}(\mathbb{R}^d)$ extend to operators in $\mathcal{M}(\mathbb{R}^d)$ we consider an operator $M = M_T \in M(\mathbb{R}^d)$, $T \in \mathcal{E}'(\mathbb{R}^d)$. We may assume that $T = (-1)^{|\beta|} t^{(\beta)}$ where $t \in L_1(\mathbb{R}^d)$ with compact support.

Here we assume that $f \in C^{\infty}(\mathbb{R}^d)$ and $\varphi \in \mathcal{D}(\mathbb{R}^d)$. Then the function

$$t(x) \varphi(y) y^{\beta} f^{(\beta)}(xy), \quad x, y \in \mathbb{R}^d$$

is in $L_1(\mathbb{R}^d \times \mathbb{R}^d)$. We use the same coordinate transformation as above. Then the function

$$\frac{\sigma(\xi)}{\xi_1 \cdots \xi_d} t\left(\frac{1}{\xi}\right) \varphi(\xi \eta) (\xi \eta)^{\beta} f^{(\beta)}(\eta), \quad \xi, \eta \in \mathbb{R}^d$$

is again in $L_1(\mathbb{R}^d \times \mathbb{R}^d)$. By Fubini's theorem and the change of variables we obtain:

$$\begin{aligned} \int \varphi(y) \left\{ \int t(x) y^{\beta} f^{(\beta)}(xy) dx \right\} dy &= \int \int t(x) \varphi(y) y^{\beta} f^{(\beta)}(xy) dx dy \\ &= \int \int \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d} t\left(\frac{1}{\xi}\right) \varphi(\xi \eta) (\xi \eta)^{\beta} f^{(\beta)}(\eta) d\xi d\eta \\ &= \int \left(\int \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d} t\left(\frac{1}{\xi}\right) \xi^{\beta} \varphi(\xi \eta) d\xi \right) (P(\theta)f)(\eta) d\eta \\ &= \int (P(\theta)f)(\eta) S_{\xi} \varphi(\xi \eta) d\eta \end{aligned}$$

where $P(\theta)f(\eta) = \xi^{\beta} f^{(\beta)}(\eta)$ and the distribution S is defined by

$$S = \frac{\sigma(\xi)}{\xi_1 \cdots \xi_d} \xi^{\beta} t\left(\frac{1}{\xi}\right).$$

Our problem now comes down to the question: when is $S \in \mathcal{D}'_H(\mathbb{R}^d)$? Then we have

$$M_T = L_S \circ f(\theta).$$

Since clearly $\text{supp } S \subset W_{\varepsilon}$ for some $\varepsilon > 0$, the question is: when does $\eta \mapsto S_{\xi} \varphi(\xi \eta)$, $\eta \in \mathbb{R}^d_{*}$ extend to a function in $C^{\infty}(\mathbb{R}^d)$?

Sufficient for that is if $\frac{\xi^\gamma}{\xi_1 \dots \xi_d} t\left(\frac{1}{\xi}\right) \in L_1(\mathbb{R}^d)$ for all $\gamma \in \mathbb{N}_0^d$ and this is equivalent to

$$x^{-\gamma} t(x) \in L_1(\mathbb{R}^d) \text{ for all } \gamma \in \mathbb{N}_0^d.$$

We have shown:

Theorem 5.3. *If $T \in \mathcal{E}'(\mathbb{R}^d)$ and there are finitely many functions $t_\beta \in L_1(\mathbb{R}^d)$ with compact support such that $T = \sum_\beta t_\beta^{(\beta)}$ and $x^{-\gamma} t_\beta(x) \in L_1(\mathbb{R}^d)$ for all $\gamma \in \mathbb{N}_0^d$ then $M_T : C^\infty(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}^d)$ extends to a map in $\mathcal{M}(\mathbb{R}^d)$.*

A condition about the behaviour of T at 0 is necessary as we have seen on Proposition 5.2.

That in the assumptions for Theorem 5.3 we need a strong vanishing condition at zero and that a condition in the spirit of the $\mathcal{O}'_H(\mathbb{R}^d)$ -condition is not sufficient is shown by the following easy example.

Example 5.4. For any $p \in \mathbb{N}_0$ we define a function t_p as follows: $t_p(x) = 0$ for $x < 0$, $t_p(x) = \chi(x)x^p/p!$ for $x \geq 0$ where $\chi \in \mathcal{D}(\mathbb{R}^d)$ and $\chi \equiv 1$ in a neighborhood of 0. Then $\delta = t_p^{(p)} + t_0$ where $t_0 \in \mathcal{D}(]0, +\infty[)$. So for any P the distribution δ is a sum of derivatives of functions with zeroes of order p in 0, but M_δ does not extend to an operator in $\mathcal{M}(\mathbb{R}^d)$ (see Proposition 5.2).

REFERENCES

- [1] P. Domański, M. Langenbruch, Representation of multipliers on spaces of real analytic functions, *Analysis* **32** (2012), 137–162.
- [2] P. Domański, M. Langenbruch, Algebra of multipliers on the space of real analytic functions of one variable, *Studia Math.* **212** (2012), 155–171.
- [3] P. Domański, M. Langenbruch, Hadamard multipliers on spaces of real analytic functions, *Adv. Math.* **240** (2013), 575–612.
- [4] P. Domański, M. Langenbruch, D. Vogt, Hadamard type operators on spaces of real analytic functions in several variables, *J. Funct. Anal.* **269** (2015), 3868–3913.
- [5] J. J. Duistermaat, J. A. C. Kolk, *Distributions. Theory and Applications*, Birkhäuser, Boston 2010.
- [6] L. Schwartz, *Théorie des distributions I*, Hermann, Paris 1957.
- [7] L. Schwartz, *Théorie des distributions II*, Hermann, Paris 1959.
- [8] R. Meise, D. Vogt: *Introduction to functional analysis*, Clarendon Press, Oxford, (1997).

- [9] D. Vogt, Operators of Hadamard type on spaces of smooth functions, *Math. Nachr.*, **288** (2015), 353–361.
- [10] D. Vogt, \mathcal{E}' as an algebra by multiplicative convolution, *arXiv: 1509.05759* (2015).