# Integer and Nonlinear Optimization Exercise 4

## Problem 1

Prove that the following statements are equivalent (Lemma 3.2 of the lecture):

- (i)  $\underline{x}^1, \dots, \underline{x}^k \in \mathbb{R}^n$  are affinely independent in  $\mathbb{R}^n$ .
- (ii)  $\underline{x}^2 \underline{x}^1, \underline{x}^3 \underline{x}^1, \dots, \underline{x}^k \underline{x}^1 \in \mathbb{R}^n$  are linearly independent in  $\mathbb{R}^n$ .
- (iii)  $(\underline{x}^1, -1), (\underline{x}^2, -1), \dots, (\underline{x}^k, -1) \in \mathbb{R}^{n+1}$  are linearly independent in  $\mathbb{R}^{n+1}$ .

#### Problem 2

Let  $P = \{\underline{x} \in \mathbb{R}^n : A\underline{x} \leq \underline{b}\}$ . Show that

- (a) If  $P \neq \emptyset$ , then P has an inner point.
- (b) If a polyhedron P is full-dimensional, then it has an interior point.

### **Problem 3** (Farkas' Lemma)

Prove the following variations of Farkas' Lemma:

- $\begin{array}{llll} \text{(a)} & \{\underline{x} \in \mathbb{R}^n : A\underline{x} = \underline{b}, \underline{x} \geq \underline{0}\} \neq \emptyset & \dot{\vee} & \{\underline{y} \in \mathbb{R}^m : \underline{y}A \geq \underline{0}, \underline{y}\,\underline{b} < 0\} \neq \emptyset. \\ \text{(b)} & \{\underline{x} \in \mathbb{R}^n : A\underline{x} = \underline{b}, \underline{x} \geq \underline{0}\} \neq \emptyset & \dot{\vee} & \{\underline{y} \in \mathbb{R}^m : \underline{y}A \leq \underline{0}, \underline{y}\,\underline{b} > 0\} \neq \emptyset. \\ \text{(c)} & \{\underline{x} \in \mathbb{R}^n : A\underline{x} \leq \underline{b}\} \neq \emptyset & \dot{\vee} & \{\underline{y} \in \mathbb{R}^m : \underline{y}A = \underline{0}, \underline{y}\,\underline{b} < 0, \underline{y} \geq \underline{0}\} \neq \emptyset. \end{array}$

#### Problem 4

Let  $S = \{\underline{s}^1, \dots, \underline{s}^k\} \subseteq \mathbb{R}^n$  be a finite set of points and let  $\underline{y} \in \mathbb{R}^n \setminus \text{conv}(S)$ . Use Farkas' Lemma to prove that there exists an inequality  $\underline{\pi} \underline{x} \leq \underline{\pi}_0$  that separates y from conv(S), that

